

Are There Ethnic Differences in Deletions in the Dystrophin Gene?

Minakshi Banerjee and Ishwar C. Verma*

Genetics Unit, Department of Pediatrics, World Health Organization Collaborating Centre in Genetics,
All India Institute of Medical Sciences, New Delhi, India

We studied 160 cases of Duchenne muscular dystrophy (DMD) drawn from all parts of India, using multiplex PCR of 27 exons. Of these, 103 (64.4%) showed intragenic deletions. Most (69.7%) of the deletions involved exons 45–51. The phenotype of cases with deletion of single exons did not differ significantly from those with deletion of multiple exons. The distribution of deletions in studies from different countries was variable, but this was accounted for either by the small number of cases studied, or by fewer exons analyzed. It is concluded that there is likely to be no ethnic difference with respect to deletions in the DMD gene. Am. J. Med. Genet. 68:152–157, 1997 © 1997 Wiley-Liss, Inc.

KEY WORDS: Duchenne muscular dystrophy; multiplex PCR; exonal deletions; ethnic differences

INTRODUCTION

The incidence of Duchenne muscular dystrophy (DMD) in different countries varies from 1:2,564–1:10,309 among male births (mean, 1:4,854), while prevalence ranges from $19.5\text{--}95 \times 10^{-6}$ (mean, 65.4×10^{-6}) in the male population [Emery, 1993]. There are no studies on the incidence of DMD in India, though Roddie and Bunday [1992] reported a higher incidence (1:1,388) of the disorder among Asian Indians in Birmingham, UK, as compared to whites, Negroes, and Pakistanis (overall incidence, 1:3,761). The prevalence of DMD in India was reported to be 47.8×10^{-6} among 20,914 males in rural Andhra Pradesh [Gouri-Devi et al., 1987]. Using the above prevalence figure and the mean worldwide incidence derived by Emery [1993], it is estimated that there are about 22,000 cases of DMD in India, while 2,687 new cases of

DMD are born every year, based on recent demographic data of India [UNICEF, 1996].

The size of the DMD gene is about 2300 kb, and it has 79 exons [Bebchuk et al., 1993; Roberts et al., 1993]. The large gene size and the 5% intragenic recombination rate [Hu et al., 1988] have made diagnostic analysis complex and laborious with standard methods using DNA probes. Deletion screening with multiplex PCR offers advantages over Southern blotting, and has replaced it [Chamberlain et al., 1988, 1992].

Most mutations in the DMD gene are intragenic deletions (65%) or duplications (5%) [Koenig et al., 1988; Forrest et al., 1988; Hu et al., 1988]. A number of studies have documented the frequency and distribution of deletions in different parts of the world [Den Dunnen et al., 1989; Gillard et al., 1989; Chamberlain et al., 1992; Covone et al., 1992; Imoto et al., 1993], and some authors have suggested that these vary with ethnic origin [Sugino et al., 1989; Tsukamoto et al., 1991; Kádasi et al., 1991; Baranov et al., 1993; Ballo et al., 1994]. There have so far been only two small studies of exonal deletions among patients with DMD in India [Sinha et al., 1992; Saxena and Verma, unpublished results]. The first study involved 5 patients who were tested for two exons, while the second study was by our own group involving 70 patients who were tested for 14 exons (data not included in present study).

We set out to investigate the frequency and pattern of intragenic deletions of the dystrophin gene by multiplex PCR, and to examine whether there are any differences among Indian subjects with DMD as compared to those in other ethnic groups.

MATERIALS AND METHODS

Patients

We evaluated 160 unrelated male patients with DMD from all parts of India. Diagnosis was established by the characteristic clinical distribution of hypertrophy and atrophy of muscles, the markedly raised creatine kinase values, and electromyography studies in all cases. Muscle biopsy, including dystrophin staining, was performed in about 20% of cases. Age range of the patients was 3–20 years, with a mean age of 8 ± 3.06 years.

DNA Extraction and PCR Primers

High molecular weight genomic DNA was isolated from peripheral blood leucocytes by the phenol-chloroform extraction method [Sambrook et al., 1989].

Contract grant sponsor: Department of Biotechnology, Government of India

*Correspondence to: Professor I.C. Verma, Genetic Unit, Department of Pediatrics, W.H.O. Collaborating Centre in Genetics, Old O.T. Building, All India Institute of Medical Sciences, New Delhi 110029, India.

Received 25 August 1995; Accepted 5 June 1996

PCR amplification of 27 exons was carried out: Pm (1), 3, 4, 6, 8, 12, 13, 17, 19, 20, 21, 22, 34, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 55, and 60. Sequences of the primers for amplification of exons Pm, 4, 8, 12, 13, 17, 19, 43, 44, 45, 48, 50, 51, and 52 were from Chamberlain et al. [1988], for exons 3, 6, 47, 49, and 60 from Beggs et al. [1990], for exons 42 and 53 from Abbs et al. [1991], for exon 22 from Bebchuk et al. [1993], and for exons 21, 34, 46, and 55 from Niemann-Seyde et al. [1992].

PCR Conditions

Amplification was carried out in six sets of reactions (Fig. 1 a,b). PCR was performed in 25- μ l volumes containing 250 ng template DNA, 12.5 pmol of each primer, 1 \times *Taq* polymerase buffer (67 mmol/l Tris-HCl, pH 8.8, 16.6 mmol/l $(\text{NH}_4)_2\text{SO}_4$, 6.7 mmol/l MgCl_2 , 6.7 mmol/l EDTA, pH 8.0, 0.5 mmol/l dNTPs (Boehringer-Mannheim, Mannheim, Germany), and 2.5 units of *Taq* DNA polymerase (Bangalore Genei Private Ltd., Bangalore, India). The cycling conditions were the same for all reaction tubes: 94°C for 7 min; followed by 25 cycles of 94°C for 30 sec, 52°C for 30 sec, and 65°C extension for 4 min, with a final extension of 65°C for 7 min, run in an MJ Research Thermocycler (Watertown, MA, USA). PCR products were observed on 4% ethidium bromide-stained Nu Sieve gels.

RESULTS

Of 160 DMD patients studied, intragenic deletions were demonstrated in the dystrophin gene in 103 (64.4%) cases. Figure 1 a,b shows representative gels.

The distribution of exonal deletions is shown in Figure 2, and the percentage frequency of different exonal deletion is depicted in Table I. Altogether 117 (42.1%) of the 278 deleted exons were spread over the region involving exons 45–51, a commonly designated “hot spot” region for deletions.

Of 103 DMD patients with deletions, 62 (60.2%) subjects had deletion of a single exon and 41 (39.8%) had deletions of multiple exons. Of the 62 single exons deleted, the commonest exons involved were numbers 50 (16.1%), 45 (14.5%), 48 (14.5%), and 51 (14.5%). The commonest exons involved in multiple deletions were numbers 48 (5.3%), 49 (13.0%), 47 (11.1%), and 50 (9.7%).

The difference in phenotype according to whether the deletion involved a single exon or multiple exons was examined (Table II). The functional status of each boy was classified according to Vignos grading [Archibald and Vignos, 1959] with special reference to the abilities to run or walk, and to confinement in the wheelchair. A comparison of the distribution of subjects with deletions of single exons as compared with those with mul-




Fig. 1. **a,b:** Six different sets of reactions, amplifying 27 exons, were used for multiplex PCR analysis for DMD. In each reaction, lane 1 represents control sample from a normal male, while the other lanes are from patients. Absence of the band indicates deletion of the specific exon.




Fig. 2. Deletions of different exons as detected by PCR are displayed by one horizontal line per patient. Majority of deletions involved exons 45–51.

TABLE I. Frequency Distribution of Deletions in Different Exons

Exons	Number of deletions			%
	Single exons	Multiple exons	Total	
Pm (exon 1)				
3	3	1	4	1.4
4		3	3	1.1
6	2	3	5	1.8
8	4	6	10	3.6
12		6	6	2.2
13		6	6	2.2
17		7	7	2.5
19	2	7	9	3.2
20		2	2	0.7
21		2	2	0.7
22	1	2	3	1.1
34		2	2	0.7
42		3	3	1.1
43		3	3	1.1
44	3	3	6	2.2
45	9	19	28	10.0
46		19	19	6.8
47	4	24	28	10.0
48	9	33	42	15.1
49	2	28	30	10.8
50	10	21	31	11.2
51	9	7	16	5.8
52		7	7	2.5
53	3	2	5	1.8
55	1		1	0.4
60				
Total	62	216	278	100

multiple exons, by using the χ^2 test, did not show any significant difference ($P = 0.37$).

We attempted to establish the beginning and the end of intragenic deletions in subjects with deletions, within the limitations of the exons tested for in the present study. Based on this information and the published sequence information on exon-intron boundaries [Koenig et al., 1989], it was determined whether the particular deletion would disturb the resultant mRNA translational reading frame. This was possible in 29 subjects with deletions. This was then correlated with clinical phenotype.

Deletions of exons 45, 45–50, and 50 are out-of-frame deletions [Koenig et al., 1989]. However, of 13 patients, 11 had a clinical phenotype of DMD, and 2 patients had phenotype of Becker muscular dystrophy (BMD). Deletions of exons 48, 45–48, 48–49, and 48–51, which are in-frame deletions, were present in 16 subjects. None of these had a phenotype of BMD, 3 had an intermediate phenotype, and 13 had a phenotype of DMD (of which 4 were wheelchair-bound). Thus, the relationship of in-frame and out-of-frame deletions to mild and severe phenotypes, respectively, was not borne out in some cases.

DISCUSSION

The PCR technique offers advantages for the diagnosis of dystrophin gene abnormalities due to its ease of application, speed, and sensitivity. It has been routinely established in different laboratories for diagnosis of DMD, and Southern blotting with cDNA probes does not seem to be superior in terms of sensitivity [Cham-

TABLE II. Distribution of Subjects With Deletions of Single and Multiple Exons According to Their Functional Status*

	Ability to walk			Wheelchair-bound
	Slowly	Unassisted	Assisted	
Single exons (n = 40) ^a	20 (50%)	4 (10%)	8 (20%)	8 (20%)
Multiple exons (n = 36) ^a	14 (38.8%)	9 (25%)	7 (19.4%)	6 (16.6%)

*Functional grading of single exon deletions vs. that of multiple exon deletions ($P = 0.37$).

^aNumber of subjects on whom complete clinical data was available.

berlain et al., 1992]. In addition, the ability to multiplex-PCR many sequences simultaneously in a single tube has permitted the development of a rapid method capable of detecting about 98% of all dystrophin-gene deletions [Chamberlain et al., 1988; Beggs et al., 1990].

In the present study, using PCR amplification of 27 exons, deletions of the DMD gene were identified in 64.4% of 160 cases of DMD. Table III compares the percentage frequency of deletions in different ethnic groups in various countries.

In the case of Japanese patients, frequency of deletions [43%, Sugino et al., 1989; 43%, Tsukamoto et al., 1991] was lower than that observed in the present study and in studies in the US [Baumbach et al., 1989; Darras et al., 1988] and Europe [Hodgson et al., 1989; Gillard et al., 1989; Den Dunnen et al., 1989; Niemann-Seyde et al., 1992; Covone et al., 1992]. However, Imoto et al. [1993] reported a higher frequency (60%). Tsukamoto et al. [1991] studied only 28 patients and analyzed only nine exons. This could explain the low frequency of deletions obtained by them. Russian [Baranov et al., 1993] and Czechoslovakian [Kádasi et al., 1991] patients also showed a much lower percentage of deletions (41% and 44%, respectively) than that from other European countries [Den Dunnen et al., 1989; Niemann-Seyde et al., 1992; Covone et al., 1992]. Baranov et al. [1993] reported deletion studies in two groups of patients. In Saint Petersburg, 11 exons were examined, while in Moscow only 10 exons were analyzed. The low deletion percentage obtained is obviously due to the analysis of only a part of the gene, as not all exons recommended for standard multiplex-PCR screening were included [Chamberlain et al., 1990; Beggs et al., 1990; Abbs et al., 1991].

Soong et al. [1991] in China analyzed only 29 patients, showing deletion in 45% of them, although they used the entire cDNA probes. Likewise, Kádasi et al. [1991] studied only a few subjects ($n = 32$), and this may be the reason for obtaining a low percentage of deletions (44%). Ballo et al. [1994] reported deletions in 40% of subjects with DMD in South Africa. They screened 15 exons by PCR and cDNA clones spanning exons 1–19 and 44–60. Among whites ($n = 32$) and Indians ($n = 24$) they observed deletions in 50% of subjects, while in black subjects ($n = 41$) deletions were detected in only 24.4%. More subjects need to be examined in the different ethnic groups in South Africa before firm conclusions can be drawn. Vainzof et al. [1991] showed deletions in 40% of Brazilian patients. However, they used only two probes for

Southern blotting and also studied a small number of patients ($n = 47$). The reason for lower frequency of deletions reported by Chamberlain et al. [1992], in a multicenter study involving 745 subjects from leading laboratories in the world, might be because only 18 primers covering nine exons were used. However, they compared the accuracy of a single PCR-multiplex amplification (nine exons) with Southern analysis using 10 cDNA probes covering the full length of the gene. The multiplex-PCR analytic method detected 82% of those deletions identified by Southern blot analysis.

From Table III it is clear that, whenever cDNA probes covering the whole gene were used for Southern analysis [Darras et al., 1988; Hodgson et al., 1989; Den Dunnen et al., 1989; Gillard et al., 1989; Baumbach et al., 1989; Imoto et al., 1993], or whenever most of the primers covering the two “hot spot” regions were used for PCR amplifications [Covone et al., 1992; Niemann-Seyde et al., 1992; present study], the frequency of deletions observed was similar. Interestingly, Covone et al. [1992] used 30 primer sets, and the frequency of deletions was almost the same as that in the present study, where 27 primer sets were used for PCR amplification.

In the present study, deletions were located mainly in one “hot spot” region (exons 45–51). Since the dystrophin gene has now been shown to have 79 exons, the area which is the “hot spot” for deletions in the case of Asian Indians is clearly the middle part of the gene. Although the percentage frequency of deletions varies in different laboratories, most deletions were observed in the middle part of the gene (exons 45–51), as in the results of the present study.

Thus, a review of the published data does not support ethnic variation in the frequency of deletions in the DMD genes, or the spread of deletions over different exons. The observed differences are explainable by a smaller number of subjects having been examined, or by analysis being restricted to a lower number of exons.

In the present study, the clinical phenotype of patients with deletion of single exons did not differ significantly from those with deletion of multiple exons. This is in conformity with earlier studies [Lindlof et al., 1988; Baumbach et al., 1989]. A better correlation exists between clinical severity of the dystrophy and the effect that an intragenic deletion has on the translational reading frame of the DMD mRNA [Monaco et al., 1988]. In the present study, we did not amplify all the exons, and Southern blotting was not

TABLE III. Comparison of Frequency of Deletions in Different Ethnic Groups*

Authors and country	Exons tested	Deletions	Deletions (%) at exons 45–51	Method
Imoto et al., 1993, Japan	Entire dystrophin cDNA	53/88 (60%)	47.2 blot	Southern blot
Tsukamoto et al., 1991, Japan	cDNA probes covering exons 4, 8, 12, 17, 19, 44, 45, 48, and 51	12/28 (43%)	75.1	PCR and Southern blot
Sugino et al., 1989, Japan	Entire cDNA probes	18/42 (43%)	MCP	Southern blot
Yang et al., 1994, China	Exons 1, 3, 6, 8, 13, 17, 43, 47, 50, 52, and 60	15/41 (37%)	NM	PCR
Soong et al., 1991, China	Entire cDNA probes	13/29 (45%)	50	Southern blot
Ballo et al., 1994, South Africa	Exons 3, 4, 6, 8, 13, 19, 42, 44, 45, 48, 50–53, and 60; cDNA probes spanning exons 1–19 and 44–60	52/128 (40%)	65	PCR and Southern blot
Hodgson et al., 1989, UK	Complete cDNA probes	163/287 (57%)	NM	Southern blot
Covone et al., 1992, Italy	Brain promoter region and Pm (1), 3, 4, 6, 8, 12, 13, 16, 17, 19–22, 29, 32, 34, 41–52, and 60	96/152 (63%)	NM	Multiplex-PCR
Den Dunnen et al., 1989, Netherlands	Entire cDNA probes	115/194 (59%)	MCP	Southern blot
Kádasi et al., 1991, Czechoslovakia	cDNA probes cf 56a, cf 56b, 1–2a, 2b–3, 4–5a, 5b–7, and 8	14/32 (44%)	NM	Southern blot
Niemann-Seyde et al., 1992, Germany	Pm (1), 3, 4, 6, 8, 12, 13, 17, 19, 43–52, and 60	35/56 (62.5%)	57.5	Multiplex-PCR
Baranov et al., 1993, Russia	Pm (1), 6, 8, 12, 17, 19, 42–45, 47, 48, 50, and 52	49/119 (41%)	73.5	PCR and Southern blot
Gillard et al., 1989, Canada	Entire dystrophin cDNA	109/181 (60%)	NM	Southern blot
Chamberlain et al., 1992, US (multicenter study)	Exons 4, 8, 17, 19, 44, 45, 48, and 51	346/745 (46%)	NM	Multiplex-PCR
Baumbach et al., 1989, US	Entire cDNA probes	90/160 (56%)	69	Southern blot
Darras et al., 1988, US	Entire cDNA probes	21/32 (66%)	MCP	Southern blot
Vainzof et al., 1991, Brazil	cDNA probes cf 56a and cf 56b	19/47 (40%)	NM	Immunofluorescence
Present study	Pm (1), 3, 4, 6, 8, 12, 13, 17, 19–22, 34, 42–44, 46–53, 55, and 60	103/160 (64%)	69.7	Multiplex-PCR

*NM, not mentioned; MCP, mostly central part of gene.

done. Therefore, the in-frame/out-of-frame and clinical phenotype could be tested in only 29 subjects. The data did not confirm that in-frame deletions have a milder phenotype, while out-of-frame deletions have a severe phenotype. However, most of the patients were children, and the full evolution of phenotype had not taken place.

This has also been observed by other investigators [Forrest et al., 1987; Monaco et al., 1988; Lindlof et al., 1988; Baumbach et al., 1989]. These results suggest that factors other than deletion of a specific exon are important in determining the phenotype in this disease [Baumbach et al., 1989].

The rapidity and simplicity of multiplex-PCR methods, combined with knowledge of the distribution of exonal deletions in an adequate number of Indian patients, has enabled us to evolve a diagnostic strategy to help the large number of affected families for carrier screening and prenatal diagnosis.

ACKNOWLEDGMENTS

We are grateful to all the families and physicians who cooperated in this study. We are also thankful to Ms. Shylamma and Mr. Swaminathan for technical assistance. This study was supported by a grant from the Department of Biotechnology, Government of India.

REFERENCES

- Abbs S, Yau SC, Clark S, Mathew CG, Bobrow M (1991): A convenient multiplex PCR system for the detection of dystrophin gene deletions—A comparative analysis with cDNA hybridization shows mistyping by both methods. *J Med Genet* 28:304–311.
- Archibald KC, Vignos PJ (1959): A study of contractures in muscular dystrophy. *Arch Phys Med Rehabil* 40:150–157.
- Ballo R, Viljoen D, Beighton P (1994): Duchenne and Becker muscular dystrophy prevalence in South Africa and molecular findings in 128 persons affected. *S Afr Med J* 84:494–497.
- Baranov VS, Gorbunova VN, Malysheva OV, Artemyeva OV, Kascheeva TK, Eygrafov OV, Polyakov AV, Lebedev VM, Kuznetsova TV, Shlykova SN, Mikhailov AV, Vakhlovsky VG (1993): Dystrophin gene analysis and prenatal diagnosis of Duchenne muscular dystrophy in Russia. *Prenat Diagn* 13:323–333.
- Baumbach LL, Chamberlain JS, Ward PA, Farwell NJ, Caskey CT (1989): Molecular and clinical correlations of deletions leading to Duchenne and Becker muscular dystrophies. *Neurology* 39:465–474.
- Bebchuk KG, Bulman DE, D'Souza VN, Worton RG, Ray PN (1993): Genomic organization of exons 22 to 25 of the dystrophin gene. *Hum Mol Genet* 2:593–594.
- Beggs AH, Koenig M, Boyce FM, Kunkel LM (1990): Detection of 98% of DMD/BMD gene deletions by polymerase chain reaction. *Hum Genet* 86:45–48.
- Chamberlain JS, Gibbs RA, Ranier JE, Nguyen PN, Caskey CT (1988): Deletion screening of the Duchenne muscular dystrophy locus via multiplex DNA amplification. *Nucleic Acids Res* 16:11141–11156.
- Chamberlain JS, Gibbs RA, Ranier JE, Caskey CT (1990): Multiplex PCR for the diagnosis of Duchenne muscular dystrophy. In Innis MA, Gelfand DH, Sninsky JJ, White T (eds): "PCR Protocols: A Guide to Methods and Applications." New York: Academic Press, pp 272–281.
- Chamberlain JS, Chamberlain JR, Fenwick RG, Ward PA, Caskey CT, Dimnik LS, Bech-Hansen NT, Hoar DI, Tantravahi U, Richards S, Covone AE, Romeo G, Abbs S, Bentley DR, Bobrow M, Rysiecki G, Ray PN, Boileau C, Junien C, Boehm C, Venne VL, Fujimura FK, Spiga I, Ferrari M, Tedeschi S, Bakker E, Kneppers LJ, van Ommen GJB, Jain K, Spector E, Crandall B, Kiuru A, Savontaus ML (1992): Diagnosis of Duchenne and Becker muscular dystrophies by polymerase chain reaction—A multicenter study. *JAMA* 267:2609–2615.
- Covone AE, Caroli F, Romeo G (1992): Screening Duchenne and Becker muscular dystrophy patients for deletions in 30 exons of the dystrophin gene by three multiplex PCRs. *Am J Hum Genet* 51:675–677.
- Darras BT, Blattner P, Harper JF, Spiro AJ, Alter S, Francke U (1988): Intragenic deletions in 21 Duchenne muscular dystrophy (DMD)/Becker muscular dystrophy (BMD) families studied with the dystrophin cDNA: Location of breakpoints on Hind III and Bgl II exon-containing fragment maps, meiotic and mitotic origin of the mutations. *Am J Hum Genet* 43:620–629.
- Den Dunnen JT, Grootsholten PM, Bakker E, Blonden LAJ, Ginjaar HB, Wapenaar MC, van Paassen HMB, van Broeckhoven C, Pearson PL, van Ommen GJB (1989): Topography of the Duchenne muscular dystrophy (DMD) gene: FIGE and cDNA analysis of 194 cases reveals 115 deletions and 13 duplications. *Am J Hum Genet* 45:835–847.
- Emery AEH (1993): "Duchenne Muscular Dystrophy," 2nd edition. Oxford: Oxford University Press, pp. 150–156.
- Forrest SM, Cross GS, Speer A, Gardner-Medwin D, Burnas J, Davies KE (1987): Preferential deletion of exons in Duchenne and Becker muscular dystrophies. *Nature* 329:638–640.
- Forrest SM, Cross GS, Flint T, Speer A, Robson KJH, Davies KE (1988): Further studies of gene deletions that cause Duchenne and Becker muscular dystrophies. *Genomics* 2:109–114.
- Gillard EF, Chamberlain JS, Murphy EG, Duff CL, Smith B, Burghes AHM, Thompson MW, Sutherland J, Oss I, Bodrug SE, Klamut HJ, Ray PN, Worton RG (1989): Molecular and phenotypic analysis of patients with deletions within the deletion-rich region of the Duchenne muscular dystrophy (DMD) gene. *Am J Hum Genet* 45:507–520.
- Gourie-Devi M, Rao VN, Prakash R (1987): Neuroepidemiological study in semiurban and rural areas in South India—Pattern of neurological disorders including motor neurone disease. In Gourie-Devi M (ed): "Motor Neurone Disease—Global Clinical Patterns and International Research." New Delhi: Oxford & IBH Publishing Co., pp 11–12.
- Hodgson S, Hart K, Abbs S, Heckmatt J, Rodillo E, Bobrow M, Dubowitz V (1989): Correlation of clinical and deletion data in Duchenne and Becker muscular dystrophy. *J Med Genet* 26:682–693.
- Hu X, Burghes AHM, Ray PN, Thompson MW, Murphy EG, Worton RG (1988): Partial gene duplication in Duchenne and Becker muscular dystrophy. *J Med Genet* 25:369–376.
- Imoto N, Arinami T, Hamano K, Matsumura K, Yamada H, Hamaguchi H, Takita H (1993): Topographic pattern of the rearrangement of the dystrophin gene in Japanese Duchenne muscular dystrophy. *Hum Genet* 92:533–536.
- Kádasi L, Géczi J, Saksová L (1991): Frequency and distribution of deletions in dystrophin gene in Duchenne muscular dystrophy patients from an East-European Slavonic population. *Gene Geograph* 5:137–140.
- Koenig M, Monaco AP, Kunkel LM (1988): The complete sequence of dystrophin predicts a rod-shaped cytoskeletal protein. *Cell* 53:219–228.
- Koenig M, Beggs AH, Moyer M, Scherpf S, Heindrich K, Bettecken T, Meng G, Savontaus M-L, Gilgenkrawtz H, Recan D, Chelly J, Kaplan J-C, Covone AE, Archidiacono N, Romeo G, Liechti-Gallati S, Schneider V, Braga S, Moser H, Darras BT, Murphy P, Francke U, Chen JD, Morgan G, Denton M, Greenberg CR, van Ommen GJB, Kunkel LM (1989): The molecular basis for Duchenne versus Becker muscular dystrophy: Correlation of severity with type of deletion. *Am J Hum Genet* 45:498–506.
- Lindlof M, Kaariainen H, van Ommen GJB, de la Chapelle A (1988): Microdeletions in patients with X-linked muscular dystrophy: Molecular-clinical correlations. *Clin Genet* 33:131–139.
- Monaco AP, Bartelson CJ, Liechti-Gallati S, Moser H, Kunkel LM (1988): An explanation for the phenotypic differences between patients bearing partial deletions of the DMD locus. *Genomics* 2:90–95.
- Murray JM, Davies KE, Harper PS, Meredith L, Mueller CR, Williamson R (1982): Linkage relationship of a cloned DNA sequence on the short arm of the X-chromosome to Duchenne muscular dystrophy. *Nature* 300:69–71.
- Niemann-Seyde S, Slomski R, Rininsland F, Ellermeyer U, Kwiatkowska J, Reiss J (1992): Molecular genetic analysis of 67 patients with Duchenne/Becker muscular dystrophy. *Hum Genet* 90:65–70.
- Roberts RG, Coffey AJ, Bobrow M, Bentley DR (1993): Exon structure of the human dystrophin gene. *Genomics* 16:536–538.
- Roddie A, Bunday S (1992): Racial incidence of Duchenne muscular dystrophy in the West Midlands. *J Med Genet* 29:280.
- Sambrook J, Fritsch EF, Maniatis T (1989): "Molecular Cloning: A Laboratory Manual." Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, pp 9.17–9.18.
- Sinha S, Pradhan S, Mittal R, Mittal B (1992): Detection of gene deletions in the central region of DMD locus using polymerase chain reaction. *Indian J Med Res* 96:297–301.
- Soong BW, Tsai TF, Su CH, Kao KP, Hsiao KJ, Su TS (1991): DNA polymorphisms and deletion analysis of the Duchenne-Becker muscular dystrophy gene in the Chinese. *Am J Med Genet* 38:593–600.
- Sugino S, Fujishita S, Kamimura N, Matsumoto T, Wapenaar MC, Deng HX, Shibuya N, Miike T, Niikawa N (1989): Molecular-genetic study of Duchenne and Becker muscular dystrophies: Deletion analysis of 45 Japanese patients and segregation analysis in their families with RFLPs based on the data from normal Japanese females. *Am J Med Genet* 34:555–561.
- Tsukamoto H, Inui K, Fukushima H, Nishigaki T, Taniike M, Tanaka J, Okada S (1991): Molecular study of Duchenne and Becker muscular dystrophies in Japanese. *J Inherited Metab Dis* 14:819–824.
- UNICEF (1996): "The State of the World's Children, 1996." Oxford, UK: Oxford University Press, p 80.
- Vainzof M, Pavanetto RCM, Pavanetto-Filho I, Rapaport D, Passos-Bueno MR, Zubrzycka-Gaarn EE, Bulman DE, Zatz M (1991): Screening of male patients with autosomal recessive Duchenne dystrophy through dystrophin and DNA studies. *Am J Med Genet* 39:38–41.
- Yang RC, Yang SL, Jin SH, Chen HW, Jong YT, Chen SS, Suzuki Y (1994): Detection of dystrophin gene deletion in Chinese Duchenne/Becker muscular dystrophy patients utilizing multiplex polymerase chain reaction. *Kaohsiung J Med Sci* 10:1–8.